

Table C-15b. Mean Ichthyoplankton Densities (No./1000 m³) at Savannah River Transects During February-July 1984.^a

River mile	American shad	Blueback herring	Striped bass	Other shad ^b	Minnows	Sunfish	Crappie	Total Ichthyoplankton ^c
LOWER FARFIELD								
29.6	1.2	0.7	0.0	2.1	1.0	8.3	5.5	24.5
40.0	1.3	0.8	0.0	2.8	0.9	9.1	8.4	30.8
50.2	1.7	0.9	0.0	3.2	2.7	12.6	4.8	32.5
60.0	1.3	2.4	0.0	4.4	1.9	6.1	10.3	33.0
69.9	0.8	0.9	0.0	3.9	3.1	3.2	5.3	24.2
79.9	1.5	2.6	0.0	4.2	2.3	1.8	4.1	25.2
89.3	3.8	1.6	0.6	4.7	4.3	2.5	3.9	30.2
97.5	6.2	4.1	0.7	11.3	3.1	4.0	5.1	44.8
110.0	10.5	1.3	1.6	1.1	5.2	4.6	3.9	35.8
120.0	5.6	1.7	4.1	2.5	5.8	3.2	3.6	34.9
NEARFIELD								
128.9	3.6	0.9	3.7	2.1	6.6	3.4	3.1	32.3
129.1	5.2	0.9	3.1	1.3	7.8	5.1	2.8	34.3
137.7	8.0	1.1	6.4	1.7	9.6	6.4	4.1	49.0
141.5	7.4	2.2	3.4	2.1	13.6	10.0	6.7	57.1
141.7	11.2	1.0	14.3	2.5	12.1	10.4	6.4	67.6
150.4	2.8	3.1	6.7	3.8	5.2	2.7	6.0	41.4
150.8	5.8	2.9	5.2	4.5	4.8	1.7	5.8	39.7
152.0	5.8	1.6	2.1	4.8	4.4	1.5	6.4	34.7
152.2	9.8	1.1	2.8	3.5	5.1	0.9	4.5	38.6
155.2	5.6	1.9	6.1	3.7	3.8	0.6	4.9	35.5
155.3 ^d	0.2	6.6	1.6	4.9	3.0	0.8	10.9	43.1
155.4	4.0	1.1	6.1	3.8	4.2	1.0	5.9	36.8
157.0	9.9	1.5	4.0	3.5	4.6	1.9	6.1	45.3
157.1 ^d	0.2	4.6	1.4	5.2	2.4	1.9	10.1	42.5
157.3	11.8	2.6	3.2	4.0	6.5	1.7	4.6	47.4
UPPER FARFIELD								
166.6	3.3	2.5	0.0	8.2	5.6	0.4	1.1	28.7
176.0	4.0	0.8	0.0	7.0	6.1	1.1	0.3	26.2
187.1	3.6	0.2	0.1	4.0	5.7	0.9	0.4	23.6

BB-3

a. Source: DOE, 1987.

b. Gizzard and/or threadfin shad.

c. Totals include taxa shown plus taxa not shown.

d. Intake canals.