

Table C-15c. Mean Ichthyoplankton Densities (No./1000 m<sup>3</sup>) at Savannah River Transects During February-July 1985.<sup>a</sup>

River mile	American shad	Blueback herring	Striped bass	Other shad <sup>b</sup>	Minnows	Sunfish	Crappie	Total Ichthyoplankton <sup>c</sup>	
LOWER FARFIELD									
89.3	47.4	0.9	2.0	2.2	2.5	1.4	0.1	67.8	
97.5	26.6	1.7	1.2	4.7	1.6	0.2	0.1	44.1	
110.0	39.6	0.6	1.5	1.2	3.2	0.2	0.1	55.6	
120.0	35.3	0.8	1.3	1.7	2.2	0.7	0.4	51.7	
NEARFIELD									
BB-3	128.9	41.7	1.4	5.3	2.2	4.1	1.0	0.2	64.2
	129.1	31.3	0.8	6.3	1.9	2.9	1.4	0.2	54.7
	137.7	34.1	1.8	0.4	2.9	1.9	1.4	0.1	55.2
	141.5	30.0	1.4	0.1	3.4	1.5	1.1	0.1	50.9
	141.7	54.6	0.9	0.5	3.7	1.8	1.8	0.3	83.2
	145.7	63.2	0.7	0.6	3.0	1.6	1.5	0.5	84.3
	150.4	22.0	1.0	2.4	2.1	0.8	0.2	0.3	44.7
	150.8	12.6	1.9	0.3	3.0	1.0	0.1	0.0	31.8
	152.0	10.6	1.4	3.7	3.2	1.6	0.1	0.1	28.7
	152.2	14.2	2.2	3.3	3.6	2.0	0.1	0.1	43.1
	155.2	18.2	2.6	16.0	3.1	1.0	0.0	0.2	67.1
	155.3 <sup>d</sup>	0.3	0.9	2.5	7.0	1.0	0.1	0.2	30.7
	155.4	19.3	0.8	13.6	4.8	1.6	0.1	0.0	57.0
	157.0	24.6	0.4	0.5	3.8	1.7	0.2	0.1	47.5
	157.1 <sup>d</sup>	0.8	1.1	0.1	3.0	0.7	0.2	0.1	22.2
	157.3	36.0	0.5	2.1	3.8	1.6	0.2	0.2	81.6
UPPER FARFIELD									
166.6	59.4	3.2	10.7	22.3	2.5	0.2	0.1	149.3	
176.0	15.7	2.4	0.1	27.2	4.5	0.4	0.2	70.9	
187.1	3.3	2.8	0.1	22.7	5.4	0.6	0.4	58.6	

a. Source: DOE, 1987.

b. Gizzard and/or threadfin shad.

c. Totals include taxa shown plus taxa not shown.

d. Intake canals.